

SAMHSA Opioid Overdose Prevention TOOLKIT:

Safety Advice for Patients & Family
Members



TABLE OF CONTENTS

Safety Advice for Patients & Family Members

Safety Advice for Patients & Family Members.....	1
What Are Opioids.....	1
Preventing Overdose	1
If You Suspect an Overdose	1
What Is Naloxone?	2
Report any Side Effects	2
Store Naloxone in a Safe Place	2
Summary: How to Avoid Opioid Overdose	2
References	3
Acknowledgments.....	4

Also see the other components of this Toolkit:

- Facts for Community Members
- Five Essential Steps for First Responders
- Information for Prescribers
- Recovering from Opioid Overdose: Resources for Overdose Survivors & Family Members

SAFETY ADVICE FOR PATIENTS & FAMILY MEMBERS

WHAT ARE OPIOIDS

Opioids include illicit drugs such as heroin and prescription medications used to treat pain such as morphine, codeine, methadone, oxycodone, hydrocodone, fentanyl, hydromorphone, and buprenorphine.

Opioids work by binding to specific receptors in the brain, spinal cord, and gastrointestinal tract. In doing so, they minimize the body's perception of pain. However, stimulating the opioid receptors or "reward centers" in the brain can also trigger other systems of the body, such as those responsible for regulating mood, breathing, and blood pressure.

A variety of effects can occur after a person takes opioids, ranging from pleasure to nausea and vomiting, from severe allergic reactions (anaphylaxis) to overdose, in which breathing and heartbeat slow or even stop.

Opioid overdose can occur when a patient misunderstands the directions for use, accidentally takes an extra dose, or deliberately misuses a prescription opioid or an illicit drug such as heroin.

Also at risk is the person who takes opioid medications pre-scribed for someone else, as is the individual who combines opioids—prescribed or illicit—with alcohol, certain other medications, and even some over-the-counter products that depress breathing, heart rate, and other functions of the central nervous system.¹

PREVENTING OVERDOSE

If you are concerned about your own use of opioids, don't wait! Talk with the health care professional(s) who prescribed the medications for you. If you are concerned about a family member or friend, urge him or her to talk to whoever prescribed the medication.

Effective treatment of opioid use disorder can reduce the risk of overdose and help a person who is misusing or addicted to opioid medications attain a healthier life. An evidence-based practice

for treating opioid addiction is the use of United States Food and Drug Administration (FDA)-approved medications, along with counseling and other supportive services. These services are available at SAMHSA-certified and DEA-registered opioid treatment programs (OTPs).^{2,3} In addition, physicians who are trained to provide treatment for opioid addiction in office-based and other settings with medications such as buprenorphine/naloxone and naltrexone may be available in your community.⁴

IF YOU SUSPECT AN OVERDOSE

An opioid overdose requires immediate medical attention. An essential first step is to get help from someone with medical expertise as soon as possible. *Call 911 immediately* if you or someone you know exhibits any of the symptoms listed below. All you have to say: "Someone is unresponsive and not breathing." Give a clear address and/or description of your location.

Signs of **OVERDOSE**, which is a life-threatening emergency, include the following:

- The face is extremely pale and/or clammy to the touch.
- The body is limp.
- Fingernails or lips have a blue or purple cast.
- The person is vomiting or making gurgling noises.
- He or she cannot be awakened from sleep or is unable to speak.
- Breathing is very slow or stopped.
- The heartbeat is very slow or stopped.

Signs of **OVERMEDICATION**, which may progress to overdose, include:

- Unusual sleepiness or drowsiness.
- Mental confusion, slurred speech, or intoxicated behavior.
- Slow or shallow breathing.
- Extremely small "pinpoint" pupils.
- Slow heartbeat or low blood pressure.
- Difficulty in being awakened from sleep.

SAFETY ADVICE FOR PATIENTS & FAMILY MEMBERS

WHAT IS NALOXONE?

Naloxone is an antidote to opioid overdose. It is an opioid antagonist that is used to reverse the effects of opioids. Naloxone works by blocking opiate receptor sites. It is not effective in treating overdoses of benzodiazepines (such as Valium®, Xanax®, or Klonopin®), barbiturates (Seconal® or Fiorinal®), clonidine, Elavil®, GHB, or ketamine. It is also not effective in treating overdoses of stimulants such as cocaine and amphetamines (including methamphetamine and Ecstasy). However, if opioids are taken in combination with other sedatives or stimulants, naloxone may be helpful.

IMPORTANT SAFETY INFORMATION. Naloxone may cause dizziness, drowsiness, or fainting.

These effects may be worse if it is taken with alcohol or certain medicines. For more information, see <http://www.fda.gov/drugs/drugsafety/postmarketdrugsafetyinformationforpatientsandproviders/ucm472923.htm>.

REPORT ANY SIDE EFFECTS

Get emergency medical help if you or someone has any signs of an allergic reaction after taking naloxone, such as hives,

difficulty breathing, or swelling of your face, lips, tongue, or throat.

Call your doctor or 911 at once if you have a serious side effect such as:

- Chest pain, or fast or irregular heartbeats.
- Dry cough, wheezing, or feeling short of breath.
- Sweating, severe nausea, or vomiting.
- Severe headache, agitation, anxiety, confusion, or ringing in your ears.
- Seizures (convulsions).
- Feeling that you might pass out.
- Slow heart rate, weak pulse, fainting, or slowed breathing.

If you are being treated for opioid use disorder (either an illicit drug like heroin or a medication prescribed for pain), you may experience the following symptoms of opioid withdrawal after taking naloxone:

- Feeling nervous, restless, or irritable.
- Body aches.
- Dizziness or weakness.
- Diarrhea, stomach pain, or mild nausea.
- Fever, chills, or goosebumps.
- Sneezing or runny nose in the absence of a cold.

This is not a complete list of side effects, and others may occur. Talk to your doctor about side effects and how to deal with them.

STORE NALOXONE IN A SAFE PLACE

Naloxone is usually handled and stored by a health care provider.

If you are using naloxone at home, store it in a locked cabinet or other space that is out of the reach of children or pets.

SUMMARY: HOW TO AVOID OPIOID OVERDOSE

1. Take medicine only if it has been prescribed to you by your doctor.
2. Do not take more medicine or take it more often than instructed.
3. Call a doctor if your pain gets worse.
4. Never mix pain medicines with alcohol, sleeping pills, or any illicit substance.
5. Store your medicine in a safe place where children or pets can- not reach it.
6. Learn the signs of overdose and how to use naloxone to keep it from becoming fatal.
7. Teach your family and friends how to respond to an overdose.
8. Dispose of unused medication properly.

READ MORE

AT <http://www.fda.gov/drugs/drugsafety/postmarketdrugsafetyinformationforpatientsandproviders/ucm472923.htm>.

REFERENCES

References

- ¹ Brunton L, Chabner B, Knollman B. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. 12th ed. New York: McGraw-Hill; 2011.
- ² Michna E, Ross EL, Hynes WL, et al. Predicting aberrant drug behavior in patients treated for chronic pain: importance of abuse history. *J Pain Symptom Manage*. 2004;28(3):250-258.
- ³ SAMHSA's National Helpline 1-800-662-HELP (4357) or 1-800-487-4889 (TDD for hearing impaired).
- ⁴ Behavioral Health Treatment Services Locator: <https://findtreatment.samhsa.gov> to search by address, city, or zip code.

ACKNOWLEDGMENTS

Acknowledgments

This publication was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by the Association of State and Territorial Health Officials, in cooperation with Public Health Research Solutions, under contract number 10-233-00100 with SAMHSA, U.S. Department of Health and Human Services (HHS). LCDR Brandon Johnson, M.B.A., served as the Government Project Officer.

Disclaimer

The views, opinions, and content expressed herein are those of the authors and do not necessarily reflect the official position of SAMHSA or HHS. Nothing in this document constitutes an indirect or direct endorsement by SAMHSA or HHS of any non-federal entity's products, services, or policies, and any reference to a non-federal entity's products, services, or policies should not be construed as such. No official support of or endorsement by SAMHSA or HHS for the opinions, resources, and medications described is intended to be or should be inferred. The information presented here in this document should not be considered medical advice and is not a substitute for individualized patient or client care and treatment decisions.

Public Domain Notice

All materials appearing in this volume except those taken directly from copyrighted sources are in the public domain and may be reproduced or copied without permission from SAMHSA or the authors. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication may be ordered from SAMHSA's Publications Ordering Web page at <http://store.samhsa.gov>. Or, please call SAMHSA at 1-877- SAMHSA-7 (1-877-726-4727) (English).

Recommended Citation

Substance Abuse and Mental Health Services Administration. SAMHSA Opioid Overdose Prevention Toolkit. HHS Publication No. (SMA) 16-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2016.

Originating Office

Division of Pharmacologic Therapies, Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 1 Choke Cherry Road, Rockville, MD 20857.
HHS Publication No. (SMA) 16-4742. First printed 2013. Revised 2014, 2016.



HHS Publication No. (SMA) 16 4742. First printed 2013. Revised 2014, 2016